

MIOSPORES FROM THE PURBECK BEDS AND MARINE UPPER JURASSIC OF SOUTHERN ENGLAND

by GEOFFREY NORRIS

ABSTRACT. Miospores from the marine Upper Jurassic and from the non-marine Purbeck Beds (Tithonian-Berriasian) of southern England are described and illustrated. Three principal palynologic suites embracing a total of 77 species (13 of which are new) are recognized, each delimiting broadly similar spore-pollen assemblages. Suite A characterizes the Upper Kimmeridgian (rotunda and pallasioides zones), the Portlandian, and the basal Lower Purbeck Beds. Suite B characterizes the remaining Lower Purbeck and the lower part of the Middle Purbeck. Suite C occupies the remaining Middle Purbeck and all of the Upper Purbeck. The Purbeck Beds of central Sussex correlate palynologically with some of the Middle and the Upper Purbeck of Dorset and with the Serpulite and Wealden 1 to 4 of north-central Germany and Holland. The lower limit of palynologic Suite C approximately coincides with a possible Jurassic-Cretaceous boundary. The successive palynologic suites represent a progressive diversification of the spore-pollen flora derived from principally gymnosperm-pteridophyte vegetation. Although filicalean spores are the most diverse elements in the assemblages as a whole, coniferalean pollen species are the most persistent as well as the numerically dominant elements. Bryophytic and cycadalean elements remain unimportant throughout.

THIS paper describes miospore assemblages from the Purbeck Beds of southern England and from the Upper Kimmeridgian and Portlandian marine sediments exposed on the Dorset coast. Couper (1958) made an extensive study of British Jurassic and Lower Cretaceous miospores but examined only a few samples from the Purbeck Beds of Durlston Bay, Dorset and Mountfield, Sussex and three palynologic assemblages from the Kimmeridgian of Scotland. He demonstrated that significant differences exist between miospore assemblages from various Jurassic and Cretaceous stages and formations but did not attempt detailed subdivision and correlation of the Upper Jurassic. Lantz (1958*b*), however, attempted a palynologic subdivision of the English Upper Jurassic in a limited study. Hughes (1958) demonstrated a relatively detailed miospore zonation of both marine and non-marine Lower Cretaceous sediments and correlated the Wealden with the marine stages by this means. Recently, Hughes and Moody-Stuart (1969) have noted a new method to correlate palynologically the Lower Cretaceous of southern England.


The present study examines the stratigraphic value of miospores for zonation and correlation of strata developed close to the Jurassic-Cretaceous boundary in southern England and for age determination of the Purbeck Beds and equivalent strata in Dorset and Sussex.

STRATIGRAPHY AND LOCATION OF SAMPLES

The stratigraphy of the sections is summarized in text-fig. 1. Location of the sections examined are indicated in text-fig. 2. Precise stratigraphic positions of the samples within these sections are given in the Appendix to Norris (1963). Reference should be made to Arkell (1947) for a detailed account of the stratigraphy and structural setting where also a full list of references to major works on the geology of the Dorset area may be found.

| | | SAMPLED INTERVAL | DORSET | SAMPLED INTERVAL | SUSSEX |
|-------------------------|--|------------------|---------------------------------|------------------|--------------------|
| WEALDEN (LOWER PART) | | | VARIEGATED MARLS AND SANDSTONES | | ASHDOWN SAND |
| | | | | | FAIRLIGHT CLAY |
| PURBECK BEDS | | | UPPER PURBECK | | "UPPER" PURBECK |
| | | | MIDDLE PURBECK | | "MIDDLE" PURBECK |
| | | | LOWER PURBECK | | "LOWER" PURBECK |
| PORTLANDIAN | | | FREESTONE SERIES | | PORTLAND SANDSTONE |
| | | | CHERTY SERIES | | |
| | | | PORTLAND SAND | | |
| KIMMERIDGIAN | | | KIMMERIDGE CLAY (UPPER PART) | | |

TEXT-FIG. 1. Generalized succession of strata close to the Jurassic-Cretaceous boundary in Dorset and Sussex. Correlations are not implied between the two areas.



TEXT-FIG. 2. Locality map of Kimmeridgian, Portlandian, and Purbeck sections examined in southern England, plus other areas mentioned in the text. Inset map shows localities on the Dorset coast within the area enclosed by a broken line on the main map.

Samples were selected on the basis of lithologies most suitable for palynomorph preservation in each section. In general, the samples from the Kimmeridge Clay are shales or silty shales and those from the Portland Sand are siltstones or fine-grained sandstones. Clastic material is almost entirely lacking in the Portland Stone and all samples from here are relatively pure limestones. Samples from the Purbeck Beds are more varied but in general grey and brown shales, silty shales, and clays yielded the best assemblages. Of the carbonate rocks in the Purbeck Beds, argillaceous limestones are the most satisfactory for palynologic examination.

TABLE I

Zonal classification of the Portlandian and upper part of the Kimmeridgian in Dorset, with thicknesses of respective rock divisions exposed at West Weare Cliff (Isle of Portland) and Hounstout (Isle of Purbeck). The strata comprising the upper two Portlandian zones constitute the Portland Stone and those of the lower two zones the Portland Sand. Numbers of samples examined are indicated.

| Stage | Zone | Isle of Portland | Rock Units | Isle of Purbeck | |
|--------------|------------------------------|------------------------------|-----------------------------------|-----------------|--|
| Portlandian | <i>Titanites titan</i> | Barren | Freestone Series 25 ft. | Not sampled | Freestone Series 50 ft. |
| | <i>Kerberites okusensis</i> | | | | |
| | <i>Glaucolithites gorei</i> | 22 samples | Portland Clay 14 ft. | 9 samples | Black Sandstones and Parallel Bands 44 ft. St. Alban's Head Marls 45 ft. White Cementstone 2 ft. Emmit Hill Marls 30 ft. Massive Bed 6 ft. |
| | <i>Zaraiskites albanii</i> | | West Weare Sandstone 40 ft. | | |
| | | | Exogyra Bed 8 ft. | | |
| | | | Upper Black Nore Beds 35 ft. | | |
| | | Black Nore Sandstone 6 ft. | | | |
| | | Lower Black Nore Beds 25 ft. | | | |
| Kimmeridgian | <i>Pavlovia pallasioides</i> | 12 samples | Kimmeridge Clay 100+ ft. | 10 samples | Hounstout Marl 50 ft. |
| | <i>Pavlovia rotunda</i> | Not sampled | (Not exposed at West Weare Cliff) | | Hounstout Clay 30 ft. Rhynchonella Marls 20 ft. Lingula Shales 40 ft. Rotunda Clays and Nodules 98 ft. Crushed Ammonoid Shales 108 ft. |

Marine Upper Jurassic

(a) *Hounstout (Isle of Purbeck)*. A complete succession from the Upper Kimmeridgian to Portlandian, as shown in Table 1, is exposed in the face of Hounstout and surrounding cliffs at Chapman's Pool, Emmit Hill, Pier Bottom, and St. Alban's Head on the Isle of Purbeck (Arkell 1935, 1947). Number of samples collected is shown in Table 1.

Arkell's (1935, 1947) zonal classification of the Kimmeridgian and Portlandian is followed, with modifications suggested by House (1958 a, b), and is also shown in Table 1. The top two ammonite zones of the Kimmeridge clay were examined in this study so that the typical shaley facies of the Kimmeridgian would be included.

(b) *West Weare Cliff (Isle of Portland)*. A continuous succession from Kimmeridge Clay to Portland Stone is exposed in the cliffs on the west side of the Isle of Portland as shown in Table 1 (Arkell 1933, 1935; House 1958*a*). The number of samples used is also shown in the Table.

Purbeck Beds

The lower boundary of the Purbeck Beds in southern England is delimited by Portlandian sediments which belong to different zones in different areas (Arkell 1953, Taitt and Kent 1958, Falcon and Kent 1960). The Purbeck Beds are terminated by the incoming of the Wealden facies (Allen 1955). Arkell (1947, 1956) used 'Purbeckian' as a stage of the Jurassic system, basing it on three ostracod zones recognizable in the Purbeck Beds. However, the Purbeckian of north-west Europe is now considered as a predominantly freshwater and continental facies developed at the top of the Jurassic and the base of the Cretaceous (Maubeuge 1962, Oertli 1963), and to avoid confusion the term will not be used.

It has been customary in British stratigraphy for more than a century to use the term 'Purbeck Beds' for all those continental, freshwater, brackish, estuarine, lagoonal or partly marine, usually fine-grained and calcareous sediments developed above the Portland Beds in southern England (Fisher 1856, Bristow 1857, Topley 1875, Woodward 1895, Arkell 1933, Howitt 1964). The desirability of using the same rock unit name for similar sediments from widely separated localities is questionable. The term Purbeck Beds is used in this study for all areas because it is still in widespread use. The erection of other formational names requires more intensive work on the detailed stratigraphy at each locality than was attempted here. The present work on the Purbeck palynologic assemblages may help to resolve some problems of the stratigraphy and correlation of the Purbeck Beds.

In a later section, reference will be made to the 'Purbeckian' of France and Switzerland developed in and around the Jura Mountains (Arkell 1956, Donze 1958). To avoid confusion and to remove any connotation of time-concordance, these sediments will be informally termed, for purposes of discussions, the 'Swiss Purbecks', 'French Purbecks', or 'Jura Purbecks', depending on the context.

Some sections of the Purbeck Beds in southern England have been zoned using ostracods. The history of the zonal classification of the Purbeck Beds has been given by Anderson (1958, 1962), the later reference with particular regard to the Upper Purbeck. Anderson (1940) proposed a more detailed subdivision of the Purbeck Beds than the original tripartite division of earlier workers. This was subsequently revised by Sylvester-Bradley (1949) and Anderson (1958). In consequence, the limits of the Lower, Middle, and Upper divisions have undergone considerable fluctuations. Anderson (1958) proposed 6 ostracod zones of which the lower 2 define the Lower Purbeck, the succeeding 3 the Middle Purbeck, and the highest zone characterizes the Upper Purbeck (Table 2). The limits of these three divisions now correspond exactly to those used by Bristow (1857).

(a) *Dorset Coast*. The type section of the Purbeck Beds is exposed on the north limb of the Purbeck anticline in Durlston Bay. There is a general attenuation of the Purbeck Beds westwards from Durlston Bay (390 ft.), Worbarrow Bay (290 ft.), Bacon Hole (250 ft.) to Lulworth Cove (179 ft.) (see Howitt 1964, fig. 7). Purbeck assemblages have

not been recovered westwards beyond Lulworth Cove. The detailed stratigraphy of the Purbeck Beds has been described by Fisher (1856), Bristow (1857), and Damon (1884). Bristow's sections have usually been taken as a basis for later work, but his rock units have been emended by Woodward (1895), Strahan (1898), and Arkell (1933). Some workers still use Bristow's rock unit nomenclature in preference to Arkell's. The two schemes are compared in Table 2, where also the ostracod zones of Anderson (1958) are indicated.

TABLE 2

Ostracod zonal classification of the Purbeck Beds from Anderson (1958), with Arkell's and Bristow's alternative stratigraphic nomenclature of the Dorset Purbeck sections; the Mountfield Purbeck is from Howitt (1964) including his suggested lithologic correlations with Dorset.

| Zones | Dorset | | Mountfield, Sussex |
|---|------------------------------|-------------------------------------|---|
| | Bristow 1857 | Arkell 1953 | |
| Upper Purbeck <i>Cypridea setina</i> | Upper Cypris Clay and Shales | Viviparus Clays | Greys Limestone Series |
| | Unio Beds | Marble Beds and Ostracod Shales | |
| | Upper Broken Shell Limestone | Unio Beds Broken Shell Limestone | |
| Middle Purbeck <i>Cypridea propunctata</i> <i>Cypridea fasciculata</i> | Chief Beef Beds | Chief Beef Beds | Shales with Beef and Clay Ironstone |
| | Corbula Beds Scallop Bed | Corbula Beds | |
| | Intermarine Beds | Upper Building Stone | Arenaceous Beds Cinder Beds Plant and Bone Beds |
| | Cinder Beds | Cinder Bed | |
| | Cherty Freshwater Beds | Lower Building Stone | |
| <i>Cypridea granulosa</i> | Marley Freshwater Beds | Mammal Bed | |
| Lower Purbeck <i>Fabarella boloniensis</i> ' <i>Cypris</i> ' <i>purbeckensis</i> | Soft Cockle Beds | Marls with Gypsum and Insect Beds | Blue Limestone Series |
| | Hard Cockle Beds | | Rounden Greys Main Gypsiferous Beds |
| | Cypris Freestone | | |
| | Broken Beds | Broken Beds | |
| Soft Cap | Caps and Dirt Beds | | |
| Hard Cap | | | |

Four Purbeck localities were examined along the Dorset Coast. The type section at Durlston Bay was examined in most detail. One hundred and twenty samples were collected from the 390-ft. section of Lower, Middle, and Upper Purbeck at this locality. Of the 85 samples treated for palynomorphs, 69 yielded assemblages.

At Worbarrow Bay, 18 samples were collected, 15 were processed, and 12 yielded palynomorphs in the 290-ft. section.

At Bacon Hole, 73 samples were collected from the 250-ft. Purbeck section. 47 samples were processed and 37 of these yielded palynomorphs.

At Lulworth Cove, 30 samples were collected from the 179-ft. section. Of these, 17 samples were processed for palynomorphs but only 5 yielded assemblages. The poor yield at Lulworth may be related to the increased proportion of carbonates at this

locality compared with areas further east. Limestone proved to be generally barren of palynomorphs. For the same reason, sections of the Purbeck Beds developed close to the edge of the basin at Teffont in the Vale of Wardour (Andrews and Jukes-Browne 1894), at Swindon, Wiltshire (Sylvester-Bradley 1941), and at Hartwell, Buckinghamshire (Ballance 1963) all proved to be barren of palynomorphs and are not considered further here.

(b) *Mountfield, Sussex*. Purbeck Beds are brought to the surface in Sussex along the central line of the Wealden dome in the Brightling anticline (Edmunds 1954). The lower part of the Purbeck Beds is known in subsurface sections in the mine of Gyproc Ltd., at Mountfield and in various boreholes. The stratigraphy, compiled from surface and subsurface sections, has been summarized by Topley (1875), White (1928), and Allen (1960*b*). A more detailed account has been given by Howitt (1964). The general succession of the Purbeck Beds in the vicinity of the mine is shown in Table 2.

At Mountfield, 31 samples were collected between the lowest gypsum seam, at the base, and the Greys Limestone Series at the top of the section, a total of almost 400 ft. (Howitt 1964). These samples were from surface outcrops and from subsurface sections in the mine and from Gypsum Mines borehole M 64 (grid reference TQ 716187). Further details of this borehole are given in Appendix 1 of Norris (1963). Of the 23 samples processed from Mountfield, 21 yielded palynomorphs.

(c) *Warlingham Borehole, Surrey*. Preliminary accounts of the succession in this borehole have been published in the *Summary of Progress of the Geological Survey* (1957, p. 29; 1958, p. 48). The Purbeck-Wealden junction has been placed at 1,892 ft. on the basis of ostracod faunas and the top of the Middle Purbeck at 1,935 ft. (Anderson 1962; Howitt 1964, p. 105). The top of the Portland Beds is recognized at 2,150 ft. but below 2,040 ft. there appears to be a repetition of lithological and faunal types in the Purbeck Beds, which is attributed to reversed faulting.

Twenty-three core samples between 1,900 and 2,027 ft. in the borehole were examined for palynomorphs. Of these, 18 yielded assemblages.

Fairlight Clay

On the coast east of Hastings, the Fairlight anticline brings the Fairlight Clay to the surface (Edmunds 1954). The Fairlight Clay is the lowest Wealden formation and is generally thought to pass laterally north and westwards into, and to be overlain by, the Ashdown Sand (Allen 1960*a*). The complete Fairlight Clay succession has been described by Topley (1875) and by White (1928), who believed that the Purbeck Beds underlie this section, close to the surface on the crest of the anticline. Allen (1955) stated that the Upper Purbeck-Wealden junction as defined by ostracods is probably situated low down in the Fairlight Clay near the base of the Wealden as traditionally defined, i.e. the base of the Fairlight Clay. A series of calcareous clays and limestones developed between 949 and 1,080 ft. in the Henfield borehole and regarded as equivalent to the basal portion of the Fairlight Clay (on ostracod faunas) '... could equally well be classed with the Purbeck...' (Taitt and Kent 1958, p. 12). The lateral change of facies at this horizon promotes the possibility that the lower part of the Fairlight Clay on the Sussex coast may be a lateral equivalent of part of the Purbeck Beds of Dorset. Allen (1960*b*, p. 7) noted 'The problem of how far the Purbeckian extends upwards (if at all) into the Fairlight facies of the Wealden remains unsolved.' A detailed palynologic

study of the entire Fairlight Clay sequence would be desirable to determine its probable equivalence to the Purbeck but was not attempted in the present study. The preliminary results of a study by Hughes and Moody-Stuart (1969) using a new correlation method, however, suggest that the Fairlight Clay is in fact a lateral facies equivalent of the upper Middle and Upper Purbeck.

EXTRACTION OF PALYNOMORPHS

Most samples examined in this study were grey or brown fine-grained clastic rocks but a few limestones were represented. These samples were prepared by first crushing about 5 gm. (20–50 gm. in the case of limestones) to pass through a 1-mm. mesh sieve. Carbonates were removed with cold dilute hydrochloric acid. Silica and silicates were dissolved in hot 50% hydrofluoric acid (40–60 minutes treatment was sufficient). By-products from this reaction are finely crystalline or in part colloidal, gelatinous white, pink, or brown precipitates usually occurring as a distinct layer or intimately mixed with the organic residue. This by-product was reported by Norem (1953) to be composed of either aluminium fluosilicates or a mixture of double fluorides of calcium, magnesium, sodium, and potassium. It interferes with subsequent preparation procedures and consequently must be removed after hydrofluoric acid treatment by repeated washing and dissolution in hot 25% hydrochloric acid.

The organic residue remaining was treated with 20 kHz ultrasound for 20–30 seconds using a generator with a 60-W power output and a magnetostrictive transducer coupled to a steel probe with a 1:1 end area ratio. The probe tip has an acoustic power output of 2.74 W/cm². Acoustic treatment was used to disperse the finely divided organic material which characterized most residues. This finely divided material was removed from each residue after ultrasonic treatment by repeated short centrifugation in water with a few drops of non-ionic detergent added (Funkhouser and Evitt 1959). Ultrasonic treatment was used before oxidation of the residues because spores and pollen are more fragile to ultrasound after oxidation (McIntyre and Norris 1964).

Most samples were not highly carbonaceous and the residues did not require more than a few minutes oxidation in concentrated nitric acid or Schulze's solution. Humic material remaining after oxidation was rare in most samples, but when present dilute ammonium hydroxide removed it effectively. Safranin-O was used to stain some palynomorph assemblages. Residues were mounted in glycerine jelly and stored in a mixture of glycerine and water with phenol added to prevent microbial attack.

Rock samples used in this study are stored in the Sedgwick Museum, Cambridge. Full locality details are listed for all samples in Appendix I of Norris (1963). Type material currently in the palynology collection of the Department of Geology, University of Toronto, will be transferred to the Sedgwick Museum for permanent storage.

SYSTEMATIC PALYNOLOGY—SPECIES LIST

The species mentioned in the following list occur in the marine Upper Jurassic and Purbeck Beds. The species are listed morphographically using Dettmann's (1963) modified scheme. Where applicable the original author's name is followed by the plate

and figure number of the illustrations given in this paper. New species and new combinations are treated thoroughly in the following section on Systematic Descriptions. The stratigraphic significance of all species is discussed in a later part of the paper.

Turma TRILETES
Suprasubturma ACAVATITRILETES
Subturma AZONOTRILETES
Infraturma LAEVIGATI

- Cyathidites anstralis* Couper 1953; Plate 102, fig. 1.
Cyathidites minor Couper 1953; Plate 102, figs. 2, 3.
Deltoidospora rafaeli Burger 1966; Plate 102, fig. 11.
Deltoidospora psilostoma Rouse 1959; Plate 102, fig. 8.
Dictyophyllidites harrisii Couper 1958; Plate 102, figs. 9, 10.
Dictyophyllidites equixinus (Couper) Dettmann 1963; Plate 102, figs. 4, 5.
Stereisporites antiquasporites (Wilson and Webster) Dettmann 1963; Plate 102, figs. 13, 14.
Concavisporites jurieusis Balme 1957; Plate 102, figs. 6, 7.
Divisisporites sp. cf. *D. euskirchenensis* Thomson and Pflug 1952; Plate 102, fig. 17.

Infraturma APICULATI

- Acauthotriletes varispinosus* Pocock 1962; Plate 102, fig. 12.
Osmundacidites wellhuanii Couper 1953; Plate 102, fig. 18.
Baculatisporites comaumeusis (Cookson) Potonié 1956; Plate 102, figs. 15, 16.
Couverrucosporites variverrucatus (Couper) comb. nov.; Plate 102, fig. 19.
Leptolepidites psarosus sp. nov.; Plate 103, figs. 2–5.
Leptolepidites epacroruatus sp. nov.; Plate 103, figs. 6–9, 11.
Rubinella major (Couper) comb. nov.; Plate 103, fig. 10.
Pilosisorites trichopapillosus (Thiergart) Delcourt and Sprumont 1955; Plate 103, fig. 1.
Pilosisorites delicatulus sp. nov.; Pl. 103, figs. 12–18; Plate 104, figs. 1, 2.

Infraturma MURORNATI

- Cicatricosisporites anstraliensis* (Cookson) Potonié 1956.
Cicatricosisporites purbeckensis sp. nov.; Plate 104, figs. 5–11.
Cicatricosisporites augicanalis Döring 1965; Plate 104, figs. 12–13; Plate 105, figs. 1, 2.
Cicatricosisporites brevilaesuratus Couper 1958; Plate 105, fig. 3.
Reticulisporites semireticulatus (Burger) comb. nov.; Plate 105, figs. 4, 5.
Lycopodiacidites cerniidites (Ross) comb. nov.; Plate 105, figs. 6, 7.
Lycopodiumsporites austroclavatidites (Cookson) Potonié 1956; Plate 105, figs. 8, 9.
Khukisporites pseudoreticulatus Couper 1958; Plate 105, fig. 11.
Microreticulatisporites diatretus sp. nov.; Plate 105, figs. 12–15.
Foveosporites canalis Balme 1957; Plate 106, fig. 3.
Tripartina sp.; Plate 105, fig. 10.

Subturma ZONOTRILETES

Infraturma AURICULATI

Trilobosporites bernissartensis (Delcourt and Sprumont) Potonié 1956; Plate 106, figs. 1, 2.

Trilobosporites apiverrucatus Couper 1958; Plate 107, figs. 9, 14.

Trilobosporites obsitus sp. nov.; Plate 106, figs. 7, 8.

Trilobosporites domitus sp. nov.; Plate 106, figs. 9, 10, 12, 13.

Appendicisporites potomacensis Brenner 1963; Plate 107, figs. 1-4, 7, 10.

Plicatella abaca (Burger) comb. nov.; Plate 106, figs. 4-6, 11; Plate 107, figs. 5, 6.

Infraturma CINGULATI

Foraminisporis wonthaggiensis (Cookson and Dettmann) Dettmann 1963; Plate 107, figs. 11, 13.

Contignisporites dorsostriatus (Bolchovitina) Dettmann 1963; Plate 107, fig. 12.

Duplexisporites problematicus (Couper) Playford and Dettmann 1965.

Infraturma TRICRASSATI

Gleicheniidites senonicus Ross 1949; Plate 107, figs. 16, 17.

Sestrosporites pseudoalveolatus (Couper) Dettmann 1963; Plate 108, fig. 5.

Coronatispora valdensis (Couper) Dettmann 1963; Plate 108, figs. 1, 2.

Suprasubturma PERINOTRILETES

Densoisporites perinatus Couper 1958; Plate 108, figs. 3, 4, 6.

Heliosporites sp.; Plate 108, figs. 7, 8, 10, 11.

Turma HILATES

Aequitriradites spinulosus (Delcourt and Sprumont) Cookson and Dettmann 1961; Plate 108, fig. 9.

Couperisporites complexus (Couper) Pocock 1962; Plate 108, fig. 13.

Januasporites tumulosus sp. nov.; Plate 108, fig. 12; Plate 109, figs. 2-4, 7.

Turma MONOLETES

Suprasubturma ACAVATOMONOLETES

Subturma AZONOMONOLETES

Infraturma SCULPTATOMONOLETI

Marattisporites scabratus Couper 1958; Plate 109, figs. 5, 6.

Anteturma POLLENITES

Turma SACCITES

Subturma MONOSACCITES

Infraturma SACCIZONATI

Cerebropollenites mesozoicus (Couper) Nilsson 1958; Plate 109, figs. 11, 12.

Subturma DISACCITES

- Alisporites bilateralis* Rouse 1959; Plate 109, figs. 14, 15.
Abietinaepollenites minimus Couper 1958; Plate 109, fig. 13.
Vitreisporites pallidus (Reissinger) Potonié 1960; Plate 109, figs. 8–10.
Podocarpidites sp. cf. *P. ellipticus* Cookson 1947; Plate 109, figs. 16, 17.
Parvisaccites radiatus Couper 1958; Plate 109, figs. 18, 19; Plate 9, fig. 1.

Subturma POLYSACCITES

- Callialasporites* sp. cf. *C. trilobatus* (Balme) Sukh Dev 1961; Plate 110, fig. 8.
Callialasporites dampieri (Balme) Sukh Dev 1961 emend.; Plate 110, figs. 2, 3.
Callialasporites obrutus sp. nov.; Plate 110, figs. 6, 7.
Callialasporites sp.; Plate 110, figs. 4, 5.

Turma ALETES

Infraturma PSILONAPITI

- Inaperturopollenites dubius* (Potonié and Venitz) Thomson and Pflug 1953; Plate 110, figs. 9, 10; Plate 111, fig. 19.
Inaperturopollenites sp.; Plate 110, figs. 11, 12.

Infraturma GRANULONAPITI

- Araucariacites australis* Cookson 1947; Plate 110, fig. 17.
Spheripollenites subgranulatus Couper 1958; Plate 110, fig. 13.

Infraturma SPINONAPITI

- Peltandripites tener* sp. nov.; Plate 110, figs. 18, 19.

Infraturma RETICULONAPITI

- Undulatasporites araneus* sp. nov.; Plate 110, figs. 14–16; Plate 111, figs. 2–10.

Turma PPLICATES

Subturma PRAECOLPATES

- Eucommiidites troedssonii* Erdtman 1948; Plate 111, figs. 13, 14, 16.
Eucommiidites minor Groot and Penny 1960; Plate 111, fig. 15.

Subturma MONOCOLPATES

- Cycadopites* sp. cf. *C. nitidus* (Balme) comb. nov.; Plate 111, figs. 11, 12.
Cycadopites carpentieri (Delcourt and Sprumont) Singh 1964; Plate 111, fig. 18.
Monosulcites sp. aff. *M. minimus* Cookson 1947; Plate 111, fig. 17.

Turma POROSES

Subturma MONOPORINES

- Exesipollenites scabrosus* sp. nov.; Plate 111, figs. 20–2.
Perinopollenites elatoides Couper 1958; Plate 112, figs. 6, 7.

MIOSPORES INCERTAE SEDIS

Classopollis torosus (Reissinger) Balme 1957; Plate 112, figs. 1–5.

Classopollis echinatus Burger 1965; Plate 112, figs. 8–13.

Classopollis hammenii Burger 1965; Plate 112, figs. 14–16; Plate 113, figs. 1–4.

Schizosporis reticulatus Cookson and Dettmann 1959; Plate 113, figs. 5, 8.

Schizosporis spriggi Cookson and Dettmann 1959; Plate 113, figs. 6, 13.

Schizosporis parvus Cookson and Dettmann 1959; Plate 113, fig. 7.

Signipollis callosus sp. nov.; Plate 113, figs. 9–12.

SYSTEMATIC DESCRIPTIONS

Turma TRILETES

Suprasubturma ACAVATITRILETES

Subturma AZONOTRILETES

Infraturma LAEVIGATI

Genus DIVISISPORITES Thomson and Pflug 1952

Divisiporites sp. cf. *D. euskirchenensis* Thomson and Pflug 1952

Plate 102, fig. 17

Description. Spores radiosymmetric, complexly trilete. Amb triangular convex to irregularly sub-circular. Laesurae long, simple, straight but dichotomizing up to the third order, reaching or almost reaching the equator. Both proximal and distal surfaces scabrate to granular. Granules usually low (occasional up to $1\ \mu$ high), closely spaced, up to $2\ \mu$ in diameter. Occasional verrucae developed at the equator up to $3\ \mu$ high and $10\ \mu$ or more in width. Exine $3\text{--}5\ \mu$ in total thickness; endexine distinct and $0.5\text{--}1\ \mu$ thick.

Dimensions (3 specimens). Equatorial diameter: $53\text{--}69\ \mu$.

Distribution. Upper Purbeck, Dorset.

Remarks. These specimens differ from *D. euskirchenensis* by the possession of a thicker exine, more complexly divided laesurae, and verrucae at the equator. Insufficient specimens were available to erect a new species. Lower Cretaceous specimens originally assigned to *D. euskirchenensis* by Cookson and Dettmann (1958) are now included in *Rouseisporites radiatus* Dettmann (1963), which is distinct in structure from the present specimens.

Infraturma APICULATI

Genus CONVERRUCOSISPORITES Potonié and Kremp 1954

Converrucosisporites variverrucatus (Couper) comb. nov.

Plate 102, fig. 19

1958 *Concavisporites variverrucatus* Couper, p. 142, pl. 22, figs. 4–5.

Remarks. This species is removed from *Concavisporites* because of the lack of curvaturae, and recombined with *Converrucosisporites* on the basis of shape and ornament.

Genus LEPTOLEPIDITES Couper 1953 emend. Norris 1968

Leptolepidites psarosus sp. nov.

Plate 103, figs. 2-5

Holotype. GN 109B/1, 40.4 124.5. Sample 59-1-6 (dark grey shale), Middle Purbeck, Chief Beef Beds, Durlston Bay (from the top of Bed 75, Bristow 1857).

Diagnosis. Spores radiosymmetric, trilete, amb rounded triangular to circular. Laesurae simple, reaching the equator, usually indistinct. Proximal face entirely granulate, or only on the contact areas. Distal face with closely spaced verrucae. Exine thin in between the projections.

Description. Laesurae straight, reaching or almost reaching the equator, occasionally quite distinct but usually difficult to see. Granules on proximal face rounded or polygonal, 1-2 μ in diameter, dense, occasionally joining up to give a sub-rugulate sculpture, rugulae about 4 μ long. When only contact faces are granular there is a distinct levigate zone separating them from the verrucate distal ornament.

Distal verrucae 3-13 μ in diameter, 1-5 μ high, rounded sub-circular, rounded polygonal, irregular or elongated. Solitary rugulae up to 17 μ long and about 5 μ wide may be interspersed among the verrucae. Verrucae closely packed forming a distinct negative reticulum with grooves about 0.5 μ wide but varying slightly. The distal verrucae may encroach to a variable extent on to the proximal face up to the contact areas.

Exine between projections less than 0.5-1 μ in thickness. Exine on proximal face less than 0.5 μ in thickness.

Dimensions. Equatorial diameter: 20-44 μ (holotype, 39 μ).

Distribution. Middle and Upper Purbeck, Dorset, Sussex, and Surrey.

Remarks. Distinguished from *Converrucosisporites proxigranulatus* Brenner (1963) by the more densely packed distal verrucae and by the presence of rugulae.

EXPLANATION OF PLATE 102

All figures $\times 750$ unless otherwise stated.

Figs. 1-3. *Cyathidites* spp. 1, *C. australis* Couper, GN 161/1, 32.0 127.4. 2-3, *C. minor* Couper, 2, GN 317/2, 44.2, 128.5. 3, GN 138/2, 58.2 123.1.

Figs. 4, 5, 9, 10. *Dictyophyllidites* spp. 4-5, *D. equixiuis* (Couper) Dettmann. 4, GN 146/1, 24.7 128.3. 5, GN 161/2, 53.0 117.7. 9-10, *D. harrisii* Couper. 9, GN 316/1, 32.6 114.9. 10, GN 316/1, 58.0 124.8.

Figs. 6, 7. *Concavisporites juriensis* Balme. 6, GN 142/2, 43.1 112.5. 7, GN 142/1, 40.1 119.6.

Figs. 8, 11. *Deltoidospora* spp. 8, *D. psilostoma* Rouse. 1113-13c, 38.3 116.9. 11, *D. rafaeli* Burger, GN 262/2, 32.2 107.5.

Fig. 12. *Acauthotriletes varispinosus* Pocock, GN 188/1, 44.0 114.5; $\times 1250$.


Figs. 13, 14. *Stereisporites antiqvasporites* (Wilson and Webster) Dettmann. 13, GN 476/1, 51.8 112.5; $\times 1250$. 14, GN 187/1, 29.5 108.6; $\times 1250$.

Figs. 15, 16. *Baculatisporites comaumensis* (Cookson) Potonié. 15, 1113-18c, 22.6 118.9. 16, GN 432/2, 46.9 104.3.

Fig. 17. *Divisisporites* sp. cf. *D. euskircheneusis* Thomson and Pflug, GN 345/1, 117.3 56.2.

Fig. 18. *Osmundacidites wellmanii* Couper, GN 316/1, 22.2 106.6.

Fig. 19. *Converrucosisporites variverrucatus* (Couper) comb. nov., GN 338/3, 50.9 124.1.



NORRIS, Late Jurassic and Purbeck miospores

Leptolepidites epacroronatus sp. nov.

Plate 103, figs. 6–9, 11

Holotype. Slide GN 147/1, 27.8 119.1. Sample 60–5–22 (dark grey-brown shale), Upper Purbeck, Marble Beds and Ostracod shales, Durlston Bay (6 ft. below the top of Bed 84, Bristow 1857).

Diagnosis. Spores radiosymmetric, trilete. Amb rounded triangular to circular. Laesurae long but not reaching equator, simple or labiate. Proximate face levigate to subgranular. Distal face verrucate with occasional echinate projections. Exine thin.

Description. Laesurae simple or with very narrow lips less than $0.75\ \mu$ wide, straight or sinuous. Ornament of proximal face very much reduced with respect to distal ornament, but distal ornament encroaches on to proximal face at apices. Distal verrucae regularly or irregularly rounded or elongate, $2\text{--}3\ \mu$ in diameter, sometimes reaching $5\ \mu$ long, up to $1\ \mu$ high and spaced about $1\ \mu$ apart. Exine $0.25\text{--}0.5\ \mu$.

Dimensions. Equatorial diameter: $12\text{--}22\ \mu$ (holotype $20\ \mu$).

Distribution. Middle and Upper Purbeck, Dorset, Sussex, and Surrey.

Genus *Rubinella* (Maljavkina 1949) Potonié 1960*Rubinella major* (Couper) comb. nov.

Plate 103, fig. 10

1958 *Leptolepidites major* Couper, p. 141, pl. 21, figs. 7–8.

Description. Spores radiosymmetric, trilete. Amb rounded triangular or occasionally almost circular. Laesurae long, simple, not reaching the equator, rather indistinct. Both proximal and distal surfaces ornamented with closely spaced or touching, almost spherical, irregularly rounded or elongated verrucae. Verrucae $2\text{--}10\ \mu$ in diameter, $1\text{--}5\ \mu$ high. Exine thickness between verrucae $1\text{--}2.5\ \mu$.

Dimensions. Equatorial diameter: $39\text{--}80\ \mu$.

Remarks. It proved impossible to split off the larger specimens as a distinct species and consequently all are included in one species with a larger size range than that indicated by Couper in his original description. The species is transferred to *Rubinella* because of the comprehensive verrucate sculpture. *Leptolepidites* is characterized by verrucae on the distal face only (Norris 1968).

Genus PILOSPORITES Delcourt and Sprumont 1955

Pilosporites delicatulus sp. nov.

Plate 103, figs. 12–18; Plate 104, figs. 1, 2

Holotype. GN 428/1, 46.0 128.4 Sample 61–7–5 (buff lignitic clay), Upper Purbeck 7 ft. above *Pahudina* Clays, Bacon Hole (middle of Bed 26 described in Norris 1963).

Diagnosis. Spores radiosymmetric, trilete. Amb triangular convex with broadly rounded apices, occasionally becoming almost circular. Laesurae short, one-third to one-half of the spore radius, straight, simple, frequently indistinct. Both proximal and distal surfaces scabrate and covered with irregularly distributed echinulate processes. Processes hair-like, very narrow, straight or curved, simply terminated or briefly bifurcate,

2–5 μ long, spaced 1–2 μ apart at the equator but up to 5 μ apart at the poles. Exine 0.5–1 μ thick.

At high magnifications (greater than 1,000 diameters) the exine is seen to be distinctly microreticulate. Lumina of reticulum less than 0.25 μ in diameter and muri also very narrow. Echinulate processes very narrow and consequently indistinct. Occasionally the bases of the processes may widen up to 0.5 μ and can be seen to be hollow.

Dimensions. Equatorial diameter: 28–40 μ (holotype 30 μ).

Distribution. Upper Purbeck of Dorset.

Remarks. This species is easily overlooked on account of its delicate ornament, or it may be confused with *Stereisporites antiquasporites* (Wilson and Webster) Dettmann.

Infraturma MURORNATI

Genus CICATRICOSISPORITES Potonié and Gelletich 1933

Cicatricosisporites purbeckensis sp. nov.

Plate 104, figs. 5–11

Holotype. Slide GN 145/1, 38.8 121.9. Sample 60–5–16 (grey, slightly calcareous shale) Upper Purbeck, Marble Beds and Ostracod Shales, Durlston Bay (Bed 84 of Bristow 1857).

Diagnosis. Spores radiosymmetric, trilete. Amb. triangular. Laesurae long, straight, simple or labiate. Proximal face levigate. Distal face with 3 or 4 triangular sets of widely spaced ribs running more or less parallel to equator. Ribs narrow and uneven in width, height, and spacing.

Description. Laesurae simple or bordered by very narrow lips about 0.5 μ wide, reaching equator. Distal ribs straight or sinuous, occasionally bifurcating, 0.25–1 μ wide, 0.5–1 μ high, spaced 0.5–2 μ apart (4 ribs and intervening lumina measure 9–12 μ). Ribs project at apices but not on sides of amb. Distal ribs encroach on to proximal face at apices. Ribs are characteristically uneven in width with swollen nodes at irregular intervals along their length. Exine about 1 μ thick, with a very thin layer of endexine distinguishable.

Dimensions. Equatorial diameter 30–48 μ (holotype 47 μ).

Distribution. Lower, Middle, and Upper Purbeck of Dorset and Sussex.

EXPLANATION OF PLATE 103


All figures $\times 750$ unless otherwise stated.

Figs. 1, 12–18. *Pilosporites* spp. 1, *P. trichopapillosus* (Thiergart) Delcourt and Sprumont, GN 338/2, 30.6 117.4. 12–18, *P. delicatulus* sp. nov. 12, proximal surface, GN 427/1, 56.6 124.9. 13–14, Holotype, median focus and distal surface respectively, GN 428/1, 46.0 128.4. 15–16, Holotype, median focus, and distal surface respectively. $\times 1,250$. 17–18, Proximal and distal surfaces respectively, GN 338/2, 119.7 22.2; $\times 1,250$.

Figs. 2–9, 11. *Leptolepidites* spp. 2–5, *L. psarosus* sp. nov. 2, Proximal surface, GN 428/1, 128.5 37.0. 3, Distal surface, GN 338/3, 36.2 1.262. 4–5, Holotype, proximal and distal surfaces respectively, GN 109B/1, 40.4 124.5. 6–9, 11, *L. epacornatus* sp. nov. 6, 8, Holotype, $\times 750$ and $\times 1,250$ respectively, GN 147/1, 27.8 119.1. 7, GN 146/2, 52.6 125.3. 9, GN 146/2, 52.7 125.5. 11, GN 190/3, 41.6 108.9.

Fig. 10. *Rubinella major* (Couper) comb. nov., GN 255/1, 41.7 119.4.

Figs. 19, 20. *Cicatricosisporites australiensis* (Cookson) Potonié. 19, GN 142/1, 54.8 108.7. 20, GN 315/3, 45.5 122.8.



NORRIS, Late Jurassic and Purbeck miospores

Remarks. Distinguished from other species of *Cicatricosisporites* by means of the narrow ribs of uneven width and spacing.

Genus RETICULISPORITES Pot. and Kremp in Weyl. and Krieger 1953

Reticulisporites semireticulatus (Burger 1966) comb. nov.

Plate 105, figs. 4, 5

1966 *Lycopodiumsporites semireticulatus* Burger, p. 247, pl. 14, fig. 4.

Remarks. The concave amb and low muri place this species in the genus *Reticulisporites*.

Genus LYCOPODIACIDITES (Couper 1953) Potonié 1956

Lycopodiacidites cerniidites (Ross 1949) comb. nov.

Plate 105, figs. 6, 7

1949 *Lycopodium cerniidites* Ross, p. 30, pl. 1, figs. 1-2.

1955 *Lycopodiumsporites cerniidites* (Ross) Delcourt and Sprumont, p. 32.

Remarks. The distal surface is rugulate, not reticulate, thus placing this species in *Lycopodiacidites*. There appears to be a morphological transition between those forms of *L. cerniidites* (Ross) with more regularly arranged regulae, to forms of *Coronatispora valdensis* (Couper) Dettmann with poorly developed circum-equatorial ridges. However the latter species has a very much greater stratigraphic and geographic distribution in the sediments examined and it is likely that each species was derived from different sources, their morphological similarity being only apparent. *Reticulatisporites pudens* Balme from the Lower Cretaceous of Western Australia is similar but much smaller and carries an imperfect reticulum on the distal face.

Genus MICRORETICULATISPORITES (Knox 1950) Potonié and Kremp 1954

Microreticulatisporites diatretus sp. nov.

Plate 105, figs. 12-15

Holotype. Slide GN 148/1, 33.2 125.6. Sample 60-5-24 (grey shaley clay), Upper Purbeck Unio Beds, Durlston Bay (Bed 82 of Bristow 1857).

Diagnosis. Spores radiosymmetric, trilete. Amb rounded triangular. Laesurae one-half to one-third of the spore radius in length, usually with narrow lips. Both proximal and distal surfaces ornamented with a perfect microreticulum with rather variable circular to polygonal lumina never exceeding $2\ \mu$ in diameter. Exine $1-2\ \mu$ thick interradially, thinning to $0.75-1\ \mu$ at apices.

Description. Amb occasionally elongated along one median. The trilete scar is rather variable in development, occasionally being simple or indistinct. Muri of the microreticulum $0.5-1\ \mu$ wide, $0.3-1.5\ \mu$ in height, of slightly variable width usually widening at the junctions. Lumina $0.5-2\ \mu$ in diameter, rounded or rounded-polygonal, varying in spacing from moderately widely spaced pits to closely spaced rounded-polygonal lumina. The lumina of the proximal reticulum may be slightly radially attenuated, particularly near the ends of the laesurae.

Dimensions. Equatorial diameter: 30–40 μ .

Distribution. Middle and Upper Purbeck, Dorset and Sussex.

Remarks. Distinguished from *Foveotriletes subtriangularis* Brenner 1963 by the microreticulate ornament on both surfaces rather than foveolate ornament primarily developed on the distal surface.

Genus TRIPARTINA Maljavkina 1949 ex Potonié 1960

Tripartina sp.

Plate 105, fig. 10

Description. Spores radiosymmetric, trilete. Amb triangular, usually concave or straight-sided, slightly undulating. Laesurae long, reaching the equator, with lips about 2 μ wide. Proximal surface unornamented. Distal surface with irregular radial grooves 0.5–2 μ wide, spaced 1–2 μ apart, occasionally anastomosing or coalescing with adjacent grooves. Both proximal and distal face may be slightly undulose. Exine 1–2 μ thick.

Dimensions. Equatorial diameter 24–46 μ .

Distribution. Middle and Upper Purbeck of Dorset, Sussex, and Surrey.

Remarks. This species is distinguished from *T. sp. cf. T. variabilis* Maljavkina described by Dettmann (1963) from the Australian Cretaceous by the more broadly rounded apices and less dense distal ornament.

Subturma ZONOTRILETES

Infraturma AURICULATI

Genus TRILOBOSPORITES Pant 1954 ex Potonié 1956

Trilobosporites obsitus sp. nov.

Plate 106, figs. 7, 8

Holotype. Slide GN 163/2, 59.2 127.5. Sample 60–5–1 (buff, silty, calcareous clay), Upper Purbeck, Unio Beds, Durlston Bay (Bed 80 of Bristow 1857).


Diagnosis. Spores radiosymmetric, trilete. Amb triangular. Laesurae long, usually simple, but rather variable. Both proximal and distal faces verrucate, but sculpture sparse and more reduced on distal face. Apices each carry a very large and prominent thickening which is circular in equatorial view and is restricted to the apical equatorial region. Exine thick.

EXPLANATION OF PLATE 104

All figures $\times 750$ unless otherwise stated.

Figs. 1, 2. *Pilosisporites delicatulus* sp. nov. 1, Median focus, GN 338/2, 122.3 23.0; $\times 1,250$. 2, Proximal surface, GN 428/2, 31.4 108.3; $\times 1,250$.

Figs. 3–13. *Cicatricosisporites* spp. 3, 4, *C. australiensis* (Cookson) Potonié, Distal and proximal surfaces respectively; GN 146/2, 32.0 111.8. 5–11, *C. purbeckensis* sp. nov. 5, Proximal surface, GN 153/2, 29.3 119.7. 6, Holotype, median focus, GN 145/1, 38.8 121.9. 7, Median focus, GN 146/2, 20.9 113.2. 8, Median focus, GN 146/2, 44.7 108.2. 9, Equatorial view, GN 153/1, 21.9 109.0. 10, Equatorial view, GN 142/2, 23.2 126.7. 11, Distal surface, GN 147/2, 32.5 125.6. 12–13, *C. angicanalis* Döring, proximal and distal surfaces respectively, GN 163/2, 40.8 122.7.



Description. Amb with roughly straight or slightly concave sides which are undulating because of verrucate ornament. Apices rounded and extended by the apical projections. Laesurae may be narrowly labiate, two-thirds or more of radius in length, occasionally reaching the equator. Proximal face with dense, rounded, or irregular verrucae, 6–12 μ in diameter, 2–3 μ high, and spaced up to 3 μ apart. Occasionally they may fuse into large indistinct rugulae. Distal ornament variable, ranging from low, indistinct, sparse irregular granules about 3 μ in diameter to rounded, low or high sparse verrucae up to 11 μ in diameter. Apical thickenings usually restricted to the equator but occasionally extending about 4 μ towards the polar areas. Exine thickness 8–10 μ over the apical thickening which is 14–20 μ wide and has a rounded aspect in equatorial views. Endexine 0.75–1.5 μ thick. Ektexine 1.5–3 μ thick between the projections.

Dimensions. Equatorial diameter 56–69 μ (holotype 63 μ).

Distribution. Upper Purbeck, Dorset.

Trilobosporites domitus sp. nov.

Plate 106, figs. 9, 10, 12, 13

Holotype. Slide GN 345/1, 57.0 111.7. Sample 60–19–3 (grey calcareous marl), Upper Purbeck, Upper Cypris Clays and Shales, Bacon Hole (6 in. above Bed 1 of Arkell 1933).

Diagnosis. Spores radiosymmetric, trilete. Amb triangular, concave. Laesurae long, straight, labiate, commissures raised. Both proximal and distal faces entirely covered in closely spaced, low, irregular granules. Exine at apices undulating and usually slightly thickened in the equatorial region here, but these thickenings not extending polewards as distinct valvae. Exine thicker on proximal than distal face.

Description. Laesurae almost reach the equator, raised 1.5–2.5 μ high at the centre and provided with narrow (1 μ) tapering lips along at least half their length. Scabrate to sub-granular ornament on both proximal and distal surfaces. Granules rather irregular in outline, very low, 0.25–1 μ in diameter and spaced not more than 0.5 μ apart. The exine becomes rather undulating around the apices tending to assume a low, poorly developed, verrucate ornament, and also is slightly thickened. Exine 2–4 μ thick, increasing at apices to between 4 and 7 μ . Exine on proximal face distinctly thicker than on distal face, about 6 μ at the centre. Endexine 0.25–0.5 μ thick. Occasionally the exine is slightly thickened interradially at the middle of the concave sides. Both proximal and distal faces convex, proximal face rather flatter than distal.

Dimensions. Equatorial diameter: 56–80 μ (holotype 65 μ).

Distribution. Upper Purbeck, Dorset.

Genus *PLICATELLA* Maljavkina 1949

Plicatella abaca (Burger) comb. nov.

Plate 106, figs. 4–6, 11; Plate 107, figs. 5, 6

1966 *Cicatricosporites abacus* Burger, p. 242, pl. 7, fig. 3.

Distribution. Middle and Upper Purbeck, Dorset, Sussex, and Surrey.

Remarks. This species is transferred to *Plicatella* on account of the apical thickenings at the equator. The distinction of *Plicatella*, characterized by weak apical thickenings, from *Appendicisporites*, with more prominent apical thickenings, is arbitrary but a convenient procedure. Hughes and Moody-Stuart (1969) have questioned the validity of the genus *Plicatella* on the basis of observations on associated spores of Cretaceous schizeaceous ferns. They have noted that occasional apical thickenings occur in populations of *Cicatricosisporites*-type spores. These thickenings may be in part the result of compression in the equatorial plane. Dispersed spores, however, are difficult to relate to associated spore populations. *Plicatella abaca* has a distinctive morphology and distribution and appears to be a discrete group. Consequently it is maintained as a distinct spore species.

Suprasubturma PERINOTRILITES
Genus HELIOSPORITES Schulz 1962
Heliosporites sp.


Plate 108, figs. 7, 8, 10, 11

Description. Spores radiosymmetric, trilete, zonate, consisting of a central body and an outer more complex layer. Amb of outer layer, excluding the zone, convex triangular with broadly rounded apices. Amb of inner body convex triangular with more sharply rounded or pointed apices. Inner body excentrically placed in relation to outer layer. Laesurae of inner body long, straight, simple, reaching the equator, occasionally indistinct. Laesurae of outer layer long, sinuous, labiate, reaching equator and frequently the outer zone. Lips 1 μ wide, slightly tapering, with fibrillar structure similar to that of outer layer. Equatorial zone 4–7 μ wide, roughly parallel to amb of outer layer, 2 μ thick and tapering very slightly towards its outer edge which is smooth except for occasional projecting spines from the dorsal surface. Entire distal surface of outer layer, including the distal surface of the zone, ornamented with irregularly distributed spines, 3–5 μ high, bases roughly circular and 3–6 μ wide, spaced 4–7 μ apart (tips spaced 8 μ apart), tapering rapidly at base but more gradually towards tips which are pointed, truncate or occasionally bifurcate. Exine of inner layer levigate, 0.25 μ or less in thickness, apparently of simple, undifferentiated structure, frequently folded, particularly near equator. Outer layer including zone, spines, and trilete mark with a fibrillar structure. Fibrils about 0.25 μ thick and anastomosing to form a ‘three-dimensional’

All figures $\times 750$.

EXPLANATION OF PLATE 105

- Figs. 1–3. *Cicatricosisporites* spp. 1–2, *C. angicanalis* Döring. 1, Proximal surface, GN 140/2, 42.6 128.5. 2, Distal surface, GN 145/2, 48.0 128.6. 3, *C. brevilaesuratus* Couper, GN 265/1, 39.8 112.9.
Figs. 4, 5. *Reticulisporites semireticulatus* (Burger) comb. nov. 4, Median focus, GN 189/2, 46.0 122.5. 5, Distal surface, GN 195/1, 21.1 110.8.
Figs. 6, 7. *Lycopodiacidites cerniidites* (Ross) comb. nov. 6, Proximal surface, GN 147/1, 29.8 128.2. 7, Distal surface, GN 421/1, 111.8 60.8.
Figs. 8, 9. *Lycopodiumsporites austroclavatidites* (Cookson) Potonié. 8, Proximal surface, GN 147/2, 43.4 128.2. 9, Distal surface, GN 154/1, 29.2 127.8.
Fig. 10. *Tripartina* sp., distal surface, GN 182/3, 26.0 101.4.
Fig. 11. *Klukisporites pseudoreticulatus* Couper, GN 148/1, 45.3 119.5.
Figs. 12–15. *Microreticulatisporites diatretus* sp. nov. 12, Holotype, proximal surface, GN 148/1, 33.2 125.6. 13, Distal surface, GN 341/2, 26.8 121.6. 14, GN 145/1 121.3 39.2. 15, Proximal surface, GN 153/2, 34.7 111.8.



NORRIS, Late Jurassic and Purbeck miospores

reticulum with lumina $0.25-1 \mu$ in diameter. Outer layer very thin on proximal surface where its fibrillar nature is difficult to determine. The outer layer is $1-2 \mu$ thick at equator and on distal surface.

Dimensions. Equatorial diameter (including zone): $30-42 \mu$.

Distribution. Upper Purbeck, Dorset.

Remarks. Insufficient material was available to erect a new species. It was considered appropriate to describe the specimens in detail, however, because they are extremely distinctive and possibly of stratigraphic importance.

These specimens are tentatively referred to *Heliosporites*. Schultz interpreted the type material of this genus as possessing a distal perispore. He did not distinguish a zone. This latter feature, together with the restriction of the spinose ornament to the distal surface, suggests affinities with the genus *Styxisporites* Cookson and Dettmann. This genus, however, does not possess a central spore body.

Considerable difficulty was encountered in interpreting the structure of these specimens owing, firstly, to the indistinct appearance of the central body; secondly, to the thinness of the proximal surface of the outer layer; and thirdly, to the fibrillar nature of this layer and the zone. All these features make elucidation of the equatorial structure, in particular, very difficult. Consequently this interpretation must be considered tentative until further specimens are available. In view of this uncertainty these specimens and the type material of *Heliosporites* may ultimately prove to have a similar structure. If this is so, these specimens are distinguished from the Lower Jurassic *Heliosporites altmarkensis* Schulz by their rather small size, thinner exine of the inner spore body, and rather shorter distal spines which are not truncated.

Turma HILATES

Genus JANUASPORITES (Pocock 1962) Singh 1964

Januasporites tumulosus sp. nov.

Plate 108, fig. 12; Plate 109, figs. 2-4, 7

Holotype. Slide GN 421/1, 55.5 123.3. Sample 61-6-2 (grey, clayey shale), Upper Purbeck, *Paludina* Clays, Lulworth Cove, Dorset (from 2 ft. 6 in. below the highest *Viviparus* limestone).

Diagnosis. Spores radiosymmetric, probably trilete, with a distinct central body. Amb rounded triangular to circular or oval. Inner body thicker and ornamented with a perfect polygonal microreticulum and usually carrying a large circular aperture on one face. Outer layer is very thin and loosely fitting around the antapical face, projecting beyond the central body at the amb and ornamented with irregularly distributed granules which are distinctly raised on hollow protuberances. Exine layers usually very thin and of indeterminate thickness.

Description. Exine is composed of two distinct layers, an inner thicker and an outer, thinner membranous layer almost completely enclosing the inner body. The inner layer bears an irregular but perfect polygonal micro-reticulum; muri $0.25-0.75 \mu$ wide; lumina $1-3 \mu$ wide and of polygonal shape but occasionally irregularly rounded. Reticulum always present at centre of antapertural face but usually becoming indistinct towards edges, sometimes showing a trilete distribution by development of stronger,

elongated lumina along 3 rays. The outer layer is very thin and is covered with rounded granules about 1–2 μ high and varying from 0.5–2 μ in diameter. Granules are distinctly raised on protuberances of the outer layer giving a characteristic L–O–L pattern in surface view. Granules spaced 1–2 μ apart and irregularly distributed. Occasional gemmate or papillate projections may be distributed amongst the granules. Outer layer loosely lies over the antapertural face and projects 1–6 μ from the amb. It occasionally encroaches up to the edges of the aperture but usually becomes indistinct on the apertural face.

Inner layer about 0.25 μ thick but usually indeterminate; outer layer of exine very thin with no distinguishable optical section visible under oil immersion, excluding the thickened granules.

Dimensions. Maximum equatorial diameter: 47–62 μ (holotype 47 μ). Minimum equatorial diameter: 32–52 μ (holotype 33 μ).

Distribution. Middle and Upper Purbeck, Dorset.

Remarks. The outer layer of *Januasporites* is not referred to in the description as either a perine or a saccus since the stratification of the inner layer is not visible. This outer layer is only loosely attached on the antapertural face but becomes closely attached to the apertural face at or just beyond the amb.

Anteturma POLLENITES

Turma SACCITES

Subturma DISACCITES

Genus PODOCARPIDITES (Cookson 1947) ex Couper 1953

Podocarpidites sp. cf. *P. ellipticus* Cookson 1947

Plate 109, figs. 16, 17

Distribution. Kimmeridgian and Portlandian of Dorset. Lower, Middle, and Upper Purbeck of Dorset, Sussex, and Surrey.

Subturma POLYSACCITES

Genus CALLIALASPORITES (Sukh Dev 1961) Potonié 1966

Callialasporites sp. cf. *C. trilobatus* (Balme 1957) Sukh Dev 1961

Plate 110, fig. 8

Description. Spores radiosymmetric, alete, with a distinct central body. Amb of central body rounded triangular, surrounded by a distinctly trilobed equatorial saccus constricted


All figures $\times 750$.

EXPLANATION OF PLATE 106

Figs. 1–2, 7–10, 12–13. *Trilobosporites* spp. 1–2, *T. bernissartensis* (Delcourt and Sprumont) Potonié-1, Median focus, GN 345/1, 47.3 127.4. 2, Distal surface, GN 163/2, 51.0 122.9. 7–8, *T. obsitus* sp. nov. 7, Holotype, median focus, GN 163/2, 59.2 127.5. 8, Median focus, GN 163/2, 24.2 114.7. 9–10, 12–13, *T. domitus* sp. nov. 9, Holotype, proximal surface, GN 345/1, 57.0 111.7. 10, Median focus, GN 345/1, 57.4 120.6. 12, Equatorial view, GN 345/1, 51.6 116.5. 13, Median focus, GN 345/1, 38.4 127.8.

Fig. 3. *Foveosporites canalis* Balme, GN 168/1, 107.6 22.3.

Figs. 4–6, 11. *Plicatella abaca* (Burger) comb. nov. 4, Median focus, GN 145/2, 49.0 12.49. 5, Median focus, GN 212/2, 45.9 126.1. 6, GN 261/1, 46.6 116.0. 11, Equatorial view, GN 212/2, 40.4 126.4.



NORRIS, Late Jurassic and Purbeck miospores

at the apices of the central body. Exine scabrate or indistinctly wrinkled. Saccus 8–10 μ wide. Exine of saccus and central body indistinct and thin.

Dimensions. Equatorial diameter 50–51 μ .

Distribution. Middle and Upper Purbeck, Dorset and Sussex.

Remarks. These grains differ from *C. trilobatus* (Balme) Sukh Dev in the scabrate rather than rugulate central body and in the overall smaller size.

Callialasporites dampieri (Balme 1957) Sukh Dev emend.

Plate 110, figs. 2, 3

- 1937 *Nelumbium* type Simpson, p. 673, fig. 2a.
 1957 *Zonalapollenites dampieri* Balme, p. 32, pl. 8, figs. 88–90.
 1958 *Zonalapollenites* cf. *trilobatus* Balme; Hughes and Couper, p. 1482, fig. 1 (e).
 1958 *Zonalapollenites dampieri* Balme (partim); Lantz, p. 925, pl. 3, fig. 34.
 1958 *Zonalapollenites trilobatus* Balme (partim); Lantz, p. 925, pl. 4, fig. 37.
 1961 *Callialasporites dampieri* (Balme) Sukh Dev, p. 48.
 1962 *Pflugipollenites dampieri* (Balme) Pocock, p. 72.

Restated diagnosis. Spores radiosymmetric, alete, with a distinct central body. Central body amb circular to rounded triangular. One face of central body distinctly convex. Equatorial saccus surrounds central body and imparts a circular outline to the entire spore. Saccus attached by a narrow area at the equator of the central body, about one-fifth to one-seventh of the total diameter in width but decreasing and becoming irregularly constricted at the apices of the central body. The saccus is never constricted so much as to completely separate into three distinct lobes. Saccus usually carries delicate radial folds which may pass into rugulate folds on the attachment area, but is never folded on the centre of the spore body. Saccus and spore body scabrate to subgranular. Saccus wall about 1 μ thick. Exine of central body 0.25–0.5 μ thick.

Description. Attachment area of saccus 2–8 μ wide, distinct or indistinct, sometimes with a wrinkled appearance due to folding. Bladder usually 8 μ wide between apices but varying from 4–13 μ . Constrictions at apices reduce bladder width to the range 1–8 μ , but these are not equal and all three are not necessarily developed on any one spore.

Dimensions. Equatorial diameter: 47–69 μ .

Distribution. Kimmeridgian, Portlandian, Lower, Middle, and Upper Purbeck of Dorset, Sussex and Surrey.

Remarks. *C. dampieri* (Balme) is similar to *C. trilobatus* (Balme). Balme (1957) noted the similarity but in his description of *C. dampieri* did not mention any constrictions of the saccus, which however, are clearly shown in his photographs (pl. 8, figs. 88–90). In the present material the sacchi are always constricted to a variable degree, frequently deeply, in three places. The more deeply constricted examples are very similar to *C. trilobatus*, particularly as Balme described the bladders of this species to occasionally coalesce to form one trilobate bladder. The central body, however, is always scabrate to subgranular rather than rugulate, the latter ornament being characteristic of *C. trilobatus*.

C. dampieri is here emended to include forms in which the bladder is constricted to a variable degree at the apices of the central body but never sufficiently deeply to clearly delimit three bladders as are found in *C. trilobatus*. It is distinguished from *C. trilobatus* by this character, by the scabrate to subgranular rather than rugulate central body, by the narrow attachment areas of the bladders to the central body, and by the restriction of the rugulate folds to this area.

Thus emended it embraces forms variously attributed to or compared with *C. dampieri* (Balme) and *C. trilobatus* (Balme) by Lantz (1958b) and Hughes and Couper (1958). Some of these trilobed, but not trisaccate forms have a clear rugulate central body (e.g. Lantz 1958, pl. 4, fig. 40) and on this character are best left in *C. trilobatus*. In his original description of *C. dampieri*, Balme noted that 'some specimens show vestigial triradiate markings' although these were not illustrated. Saccate grains with trilete scars of various types and development have been illustrated by Hughes and Couper (1958) and Lantz (1958b) and attributed to Balme's species of *Zonalapollenites*. It seems advisable to remove these from *Callialasporites dampieri* (Balme) and *C. trilobatus* (Balme) to genera of the Triletisacciti since some of the trilete marks illustrated by these authors appear to be quite well developed and not vestigial.

All figures $\times 750$.


EXPLANATION OF PLATE 107

- Figs. 1-4, 7, 10. *Appendicisporites potomacensis* Brenner. 1, 2, Median focus and distal surface respectively, GN 427/1, 48.0 117.6, 3, Proximal surface, GN 265/1, 122.3 44.1. 4, Median focus, GN 265/1, 50.2 111.0. 7, Median focus, GN 265/1, 48.2 123.4. 10, Median focus, GN 138/2, 50.0 102.6.
- Figs. 5, 6. *Plicatella abaca* (Burger) comb. nov., distal and proximal surfaces respectively, GN 145/2, 45.5 108.4.
- Fig. 8. *Duplexisporites problematicus* (Couper) Playford and Dettmann, GN 109B/1, 44.2 128.6.
- Figs. 9, 14. *Trilobosporites apiverrucatus* Couper. 9, Proximal surface, GN 338/1, 34.9 124.5. 14, Proximal surface, GN 341/1, 116.8 25.8.
- Figs. 11, 13. *Foraminisporis wouthaggiensis* (Cookson and Dettmann) Dettmann. 11, GN 428/1, 58.3 110.7. 13, GN 428/1, 44.6 114.0.
- Fig. 12. *Contignisporites dorsostriatatus* (Bolchovitina) Dettmann, GN 163/2, 32.4 119.1.
- Fig. 15. *Coronatispora valdensis* (Couper) Dettmann; Damaged specimen showing structure of distal surface, GN 154/1, 27.2 127.8.
- Figs. 16, 17. *Gleicheniidites senonicus* Ross. 16, GN 146/2, 37.4 124.4. 17, GN 147/2, 28.2 116.9.


EXPLANATION OF PLATE 108

All figures $\times 750$ unless otherwise stated.

- Figs. 1, 2. *Coronatispora valdensis* (Couper) Dettmann. 1, Proximal surface, GN 187/2, 47.4 125.3. 2, Distal surface, GN 152/1, 42.5 112.3.
- Figs. 3, 4, 6. *Densosporites perinatus* Couper. 3, Proximal surface, GN 262/1, 41.5 123.3. 4, Median focus, GN 338/1, 52.8 111.2. 6, Median focus, GN 338/3, 50.4 127.9.
- Fig. 5. *Sestrosporites pseudoalveolatus* (Couper) Dettmann, GN 257/1, 44.1 116.4.
- Figs. 7, 8, 10, 11. *Heliosporites* sp., Median foci. 7, GN 338/2, 37.1 110.7. 8, GN 427/2, 34.9 112.6. 10, GN 338/2, 44.9 117.4. 11, GN 262/1, 53.3 114.2.
- Fig. 9. *Aequitriradites spinulosus* (Cookson and Dettmann) Cookson and Dettmann, GN 428/2, 50.5 124.7.
- Fig. 12. *Januasporites tumulosus* sp. nov., holotype, distal surface; GN 421/1, 55.5 123.3; $\times 1,250$.
- Fig. 13. *Couperisporites complexus* (Couper) Pocock; distal surface, GN 265/1, 59.0 115.3.



NORRIS, Late Jurassic and Purbeck miospores



NORRIS, Late Jurassic and Purbeck miospores

Callialasporites obrutus sp. nov.

Plate 110, figs. 6, 7

Holotype. Slide GN 148/1, 48.4 109.6. Sample 60-5-24 (grey, shaley clay), Upper Purbeck, Unio Beds, Durlston Bay (from Bed 82 of Bristow 1857).

Diagnosis. Spores radiosymmetric, alete, with a distinct central body. Amb of central body circular to oval, surrounded by a saccus imparting an irregularly circular outline to the entire spore. Central body outline not clear, saccus not distinctly attached to it. Saccus very thin, standing out from central body one-tenth to one-twelfth of the total diameter, scabrate to sub-granular, carrying coarse and also fine radial folds on equatorial bladder, these passing into irregular rugulate folds on central body. Central body granular to rugulate, finer sculpture elements at poles. Exine of central body (?endoexine) thin, about 0.25μ or less in thickness. Bladders (?ektexine) also thin, about 0.25μ thick.

Description. Bladders very narrow, usually about 5μ wide but varying $4-9 \mu$. Bladder folded coarsely and approximately radially. Central body usually indistinct.

Dimensions. Equatorial diameter (including sacci): $38-69 \mu$ (holotype 61μ). Central body diameter: $29-44 \mu$ (holotype 44μ).

Distribution. Kimmeridgian, Dorset. Lower, Middle, and Upper Purbeck, Dorset, Sussex, and Surrey.

Remarks. *Callialasporites obrutus* sp. nov. is distinguished from *Zonalapollenites segmentatus* Balme by the possession of a very thin-walled central body.

Callialasporites sp.

Plate 110, figs. 4, 5

Description. Spores radiosymmetric, trilete, with a narrow equatorially attached saccus. Laesurae long, sinuous, labiate, $1-2 \mu$ wide reaching beyond amb of central body and occasionally reaching amb of the saccus when they fan out into folds. Amb of both saccus and central body circular to rounded triangular. Saccus scabrate but thrown into rugulate, rather than irregular folds where attached to central body. Saccus projects evenly beyond amb of central body a distance equal to one-sixth to one-tenth of the total diameter ($4-8 \mu$). Saccus almost unfolded or carrying rather coarse, irregular radial rugulate folds up to 2μ wide and 4μ long.

Dimensions. Equatorial diameter (including sacci): $39-64 \mu$.

Distribution. Portlandian, Dorset. Middle and Upper Purbeck, Dorset and Surrey.

Remarks. Apart from the prominent trilete mark, this species is similar to *Callialasporites obrutus* sp. nov. in possessing an equatorial, radially folded bladder which is attached to a rugulate central body.

Turma ALETES

Infraturma PSILONAPITI

Genus INAPERTUROPOLLENITES (Pflug ex Thomson and Pflug 1953) Potonié 1958

Inaperturopollenites sp.

Plate 110, figs. 11, 12

Description. Spores small, spheroidal or occasionally of irregular shape owing to folding, inaperturate. Exine relatively thick and rigid, unfolded or carrying short arcuate folds. Exine 0.25–1 μ in thickness, usually about 0.75 μ .

Dimensions. Diameter: 9–15 μ .

Distribution. Kimmeridgian and Portlandian of Dorset and Sussex. Lower, Middle, and Upper Purbeck of Dorset, Sussex, and Surrey.

Infraturma SPINONAPITI
Genus PELTANDRIPITES Wodehouse 1933
Peltandripites tener sp. nov.

Plate 110, figs. 18, 19

Holotype. Slide GN 316/1, 42.0 123.7. Sample WM 2024/2 (grey, calcareous shale), Middle Purbeck, Warlingham borehole, Surrey.

Diagnosis. Spores radiosymmetric, inaperturate, spherical, folded, entirely covered in

All figures \times 750.


EXPLANATION OF PLATE 109

- Fig. 1. *Couperisporites complexus* (Couper) Pocock; distal surface, GN 338/1, 43.4 111.0.
Figs. 2–4, 7. *Jannasporites tumulosus* sp. nov. 2, 3, Holotype, median focus and distal surface respectively; GN 421/1, 55.5 123.3. 4, Median focus, showing circular aperture; GN 345/1, 41.51 28.5. 7, Median focus, GN 138/2, 40.4 117.2.
Figs. 5, 6. *Marattisporites scabratus* Couper. 5, Polar view, GN 163/2, 51.2 111.5. 6, Equatorial view, GN 147/1, 27.8 119.0.
Figs. 8–10. *Vitreisporites pallidus* (Reissinger) Potonié. 8, GN 138/2, 28.0 122.0. 9, GN 154/2, 45.7 121.7. 10, GN 153/2, 30.7 113.5.
Figs. 11, 12. *Cerebropollenites mesozoicus* (Couper) Nilsson. 11, GN 152/1, 38.1 117.9. 12, GN 152/1, 48.0 111.1.
Fig. 13. *Abietinaepollenites minimus* Couper, GN 265/2, 46.4 111.3.
Figs. 14, 15. *Alisporites bilateralis* Rouse. 14, GN 143/2, 42.6 120.8. 15, GN 259/2, 28.6 127.2.
Figs. 16, 17. *Podocarpidites* sp. cf. *P. ellipticus* Cookson. 16, GN 431/1, 118.8 39.3. 17, GN 421/1, 47.5 123.1.
Figs. 18, 19. *Parvisaccites radiatus* Couper. 18, Oblique polar view, GN 146/2, 37.0 124.3. 19, Oblique equatorial view, GN 186/2, 49.5 112.6.


EXPLANATION OF PLATE 110

All figures \times 750 unless otherwise stated.

- Fig. 1. *Parvisaccites radiatus* Couper, equatorial view, GN 138/2, 50.8 128.2.
Figs. 2–8. *Callialasporites* spp. 2–3, *C. dampieri* (Balme) Sukh Dev. 2, 1113–13C, 44.2 119.4. 3, GN 421/1, 46.6 119.3. 4–5, *C.* sp. 4, GN 152/1, 29.5 123.2. 5, GN 152/2, 36.1 121.8. 6–7, *C. obrutus* sp. nov. 6, Holotype, GN 148/1, 48.4 109.6. 7, GN 431/1, 23.9 118.2. 8, *C.* sp. cf. *C. trilobatus* (Balme) Sukh Dev, GN 255/2, 49.7 120.2.
Figs. 9–12. *Inaperturopollenites* spp. 9–10, *I. dubius* (Potonié and Venitz) Thomson and Pflug. 9, GN 196/1, 56.8 127.7. 10, GN 345/1, 35.9 108.6. 11–12, *I.* sp. 11, GN 272/1, 44.9 117.0; \times 1,250. 12, GN 421/1, 39.1 120.4; \times 1,250.
Figs. 13. *Spheripollenites subgrammlatus* Couper, GN 482/1, 34.6 109.3.
Figs. 14–16. *Undulatasporites aranens* sp. nov. 14, 16, High and median foci respectively, GN 421/2, 43.2 126.8. 15, High focus, GN 421/2, 47.1 110.0.
Fig. 17. *Arancariacites anstralis* Cookson, GN 265/2, 46.6 119.8.
Fig. 18, 19. *Peltandripites tener* sp. nov. 18, Holotype, GN 316/1, 42.0 123.7; \times 1,250. 19, Tetrad, GN 281/1, 46.4 115.0.



NORRIS, Late Jurassic and Purbeck miospores



NORRIS, Late Jurassic and Purbeck miospores

short, closely spaced, irregularly distributed spines. Exine very thin, usually less than 0.25μ in thickness.

Description. Grains always carry many arcuate and crescentic folds. Spines $0.75\text{--}1 \mu$ long, usually about 0.25μ wide but occasionally up to 1μ wide, spaced irregularly $0.25\text{--}1 \mu$ apart. In some corroded specimens some of the spines are reduced to low granules. Rarely the grains occur in tetrads. Sometimes the grains are ruptured in an arcuate fashion. Optical section of exine indistinct, occasionally reaching about 0.25μ thick but never thicker.

Dimensions. Maximum diameter: $33\text{--}55 \mu$ (holotype 50μ).

Distribution. Kimmeridgian and Portlandian, Dorset. Lower and Upper Purbeck, Dorset.

Remarks. This species is easy to confuse with some acritarchs with short processes. It is only distinguished with difficulty but is clearly a pollen grain on account of its occurrence in tetrads.

Peltrandripites tener is distinguished from *Araucariacites australis* Cookson by its smaller size, much thinner exine, and clearly echinulate rather than granular ornament.

Infraturma RETICULONAPITI

Genus UNDULATASPORITES Leschik 1955

Undulatasporites araneus sp. nov.

Plate 110, figs. 14–16; Plate 111, figs. 2–10

Holotype. Slide GN 345/1, 25.3 107.2. Sample 60–19–3, (grey calcareous marl), Upper Purbeck, Upper Cypris Clays and Shales, Bacon Hole (6 in. above Bed 1 of Arkell 1933).

Diagnosis. Spores radiosymmetric, alete. Amb circular to oval. Exine very thin and ornamented with irregular rugulae coalescing into a very imperfect, irregular microreticulum consisting of narrow muri of constant width, and elongated irregular lumina with a radially elongated arrangement towards the periphery. Exine $0.25\text{--}1 \mu$ thick, possibly tectate.

Description. Muri and rugulae $0.25\text{--}1 \mu$ wide, of constant width along their length, 0.25μ or less up to 0.5μ high. Luminae $0.5\text{--}1 \mu$ wide and up to 3μ long, tortuous and bounded by anastomosing muri which show both angular and rounded bends in their courses. Spores sometimes show concentric folds close to the periphery.

Dimensions. Maximum equatorial diameter: $21\text{--}30 \mu$ (holotype 26μ).

Distribution. Upper Purbeck, Dorset.

Remarks. *Undulatasporites araneus* sp. nov. is distinguished from *Undulatasporites anguineus* Leschik by the overall smaller size, thinner exine, and shorter rugulae.

Turma PPLICATES

Subturma MONOCOLPATES

Genus CYCADOPITES Wodehouse 1933 ex Wilson and Webster 1946

Cycadopites sp. cf. *C. nitidus* (Balme 1957) comb. nov.

Plate 111, figs. 11, 12

1957 *Entylissa nitidus* Balme, p. 30, pl. 6, figs. 78–80.

1962 *Ginkgocycadophytus nitidus* (Balme) de Jersey, p. 12.

Distribution. Kimmeridgian, and Portlandian, Dorset. Lower, Middle, and Upper Purbeck, Dorset and Surrey.

Remarks. These grains have a slightly greater over-all size range (18–39 μ long; 13–27 μ broad) than those described by Balme but also differ in their tectate and scabrate to infrapunctate exine (rather than ‘smoothly or faintly granulate’). The exine, however, has a granular appearance in some corroded specimens. Balme (1957) believed that *C. nitidus* was derived from plants of cycadalian or bennettitalian affinities. The stratification of the exine of the present grains is not in conflict with this. Pollen grains of the modern genera *Bowenia*, *Cycas*, and *Macrozamia* belonging to the Cycadaceae and are illustrated by Erdtman (1957, figs. 10, 17, 46). All have tectate exines with rod-like elements supporting the tectum and closely resemble *C. sp. cf. C. nitidus*. The exines of some of the modern forms are distinctly crassitegellate whereas the tectum of *C. sp. cf. C. nitidus* is frequently seen to be just thinner than the gap between it and the endexine but merging into a crassitegellate exine.

Turma POROSES

Subturma MONOPORINES

Genus EXESIPOLLENITES Balme 1957

Exesipollenites scabrosus sp. nov.

Plate 111, figs. 20–2

Holotype. Slide GN 316/1, 45.5 109.5. Sample WM 2024/2 (grey calcareous shale), Purbeck Beds, Warlingham borehole.

Diagnosis. Spores radiosymmetric, monoporate. Amb rounded triangular to circular. Both proximal and distal faces rather flattened. Distal face with a circular thickening around the centre with an over-all diameter about half the equatorial diameter. At the distal pole at the centre of the thickening is a rather thinner, distinctly depressed circular

EXPLANATION OF PLATE 111

All figures $\times 1,250$ unless otherwise stated.

Fig. 1. *Peltandripites tener* sp. nov., GN 281/1, 54.5 121.1.

Figs. 2–10. *Undulatasporites araneus* sp. nov. 2–3, High and median foci respectively, GN 341/1, 27.8 128.0. 4–5, Holotype, high and median foci respectively, GN 345/1, 25.3 107.2. 6–7, GN 345/1, 30.5 112.4. 8, High focus, GN 345/1, 107.3 25.9. 9, Median focus, GN 421/1, 56.1 127.6. 10, GN 345/1, 24.3 108.9.

Figs. 11, 12. *Cycadopites* sp. cf. *C. nitidus* (Balme) comb. nov. 11, GN 338/3 49.4 119.8; $\times 750$. 12, GN 196/1 43.0 109.3; $\times 750$.


Figs. 13–16. *Eucommiidites* spp. 13–14, 16, *E. troedssonii* Erdtman, $\times 750$. 13, GN 163/2, 40.3 109.9. 14, GN 163/2, 48.7 125.9. 16, GN 148/1, 38.1 117.2. 15, *E. minor* Groot and Penny, GN 345/1, 25.9 107.4; $\times 750$.

Fig. 17. *Monosulcites* sp. aff. *M. minimus* Cookson, GN 344/1, 39.2 128.6; $\times 750$.

Fig. 18. *Cycadopites carpentieri* (Delcourt and Sprumont) Singh, GN 184/1, 41.9 108.8; $\times 750$.

Fig. 19. *Inaperturopollenites dubius* (Potonić and Venitz) Thomson and Pflug, GN 153/2, 23.2 122.1; $\times 750$.

Figs. 20–22. *Exesipollenites scabrosus* sp. nov. 20, GN 316/1, 109.3 45.3; $\times 750$. 21–2, Holotype, distal surface, $\times 750$ and $\times 1,250$ respectively; GN 316/1, 45.5 109.5.



NORRIS, Late Jurassic and Purbeck miospores

